

Population synthesis in astrophysics

S.B. Popov¹, and M.E. Prokhorov¹

¹ Sternberg Astronomical Institute, Universitetski pr. 13, 119992, Moscow, Russia,
polar@sai.msu.ru, mike@sai.msu.ru

Abstract

We briefly describe different astrophysical applications of a population synthesis method. In some details we discuss the population synthesis of close binary systems and of isolated neutron stars.

1 Introduction

In this lecture notes we give a brief introduction into *population synthesis in astrophysics*.

A *population synthesis* is a method of a direct modeling of relatively large populations of weakly interacting objects with non-trivial evolution. As a rule, the evolution of the objects is followed from their birth up to the present moment.

In the following sections we describe a necessity of this type of modeling and give a general description of the method (secs. 2 and 3). Then in some details we discuss studies of close binaries (sec. 4) and of isolated neutron stars (sec. 5). In the sixth section we give examples of other types of population synthesis studies; namely, we describe modeling of galactic spectra via simulations of stellar populations and modeling of the X-ray background radiation. Finally we present our conclusions. Since many examples are just briefly touched, we provide a list of references to important papers and reviews.

Before we start it is useful to make one comment on our definitions. There are two main approaches and both are sometimes called *population synthesis*. One has a very wide area of application. In this method calculations are started with some initial conditions, and a population is followed from its birth. Some authors call it *evolutionary synthesis* (see [1] for example). However, more often it is called *evolutionary population synthesis* (see for example [2, 3, 4]). Another method, used mainly for studies of stellar populations (see sec.6.1), is simply called *population synthesis* in [1] and in some other papers. This method is often called *empirical population synthesis* (see [2]). The second way of notation is probably better, as it allows not to mix the two approaches but to save the main part of notation – population synthesis – in both. Below we shall mainly speak about the first method (except sec. 6). If it does not mislead the reader, we shall call it just *population synthesis* (PS).

2 Necessity of population synthesis studies in astrophysics

A necessity of the PS in astrophysics is mainly determined by a specific character of astrophysics as a science. We cannot make direct experiments with objects under study: experiments are replaced by observations. An observation in a sense is just a snapshot of some object(s) with very long evolutionary time scale. It is impossible to follow an evolution of a given galaxy or a star simply observing it. Often an evolutionary path can be explored only by modeling.

Another specific feature of astronomy is that “*we are searching where it is brighter*”. I.e. sometimes we observe objects which are not typical representatives of some population, but oppositely are very peculiar. We can make an attempt to construct a theory to describe a few particular observed sources of some type. Nevertheless, it is necessary to check if this theory is consistent with an existence of other (probably unobserved yet) sources of the same type. Thus, finally it is necessary to study populations, not just single (easily observable) objects.

It is important to note that in many cases observations of individual objects are not available, only integral characteristics can be obtained. In that case for better understanding of properties of an observed population it is also necessary to model it.

The third feature we want to note is like that. Many astronomical projects (satellites, networks of telescopes, etc.) are very expensive. It is necessary to justify their construction in advance. This means that it is necessary, for example, to convince a community that new interesting, previously unobserved objects can be found. It demands an analysis of populations of unobserved sources basing on any available data on their already detected “relatives”.

Two main goals of a population synthesis modeling are the following. At first the PS can be used to test or derive some parameters of a population under consideration (for example, initial spin periods of neutron stars, an initial mass function of stars) by comparison of a real (observed) and a theoretical (modeled) populations of sources. In astronomy very often we cannot directly obtain initial and evolutionary parameters from observations in a model independent way. The PS is one of very few instruments to derive unknown properties of astronomical objects via a computer modeling. If the whole set of parameters of a population is successfully tested, i.e. an output of a model is in good correspondence with observations, then we have a quantitative explanation of the origin and the evolution of the discussed type of objects.

The *Absolute PS* of that kind might contain an artificial universe in a computer. If all

properties of the artificial and real universes coincide, we can conclude that we have build a self-consistent picture of the physical world with all initial and evolutionary parameters in correspondence with data of observations and experiments. So we can call the first approach as *testing and explanation*.

The second goal of a PS model is to *predict* properties of unobserved sources (for example, the coalescence rate of binary supermassive black holes detectable by LISA, or the rate of neutron stars coalescence detectable by LIGO). In that case uncertainties of a prediction are mainly determined by uncertainties of the input parameters. “*The reliability of a population synthesis is as good as the input assumptions*” [5]. Of course, a future comparison with new observational data will give an opportunity to test assumptions of a model.

Now let us discuss some basic features of a PS which may be applied to models of different kind.

3 General properties of population synthesis models

In our opinion the PS method as it is used in astrophysics does not have direct analogues in other fields of science. For example, most of objects studied in physics (elementary particles, atoms, molecules, etc.) are much simpler than astronomical objects and have much more trivial evolution. On the other hand, behaviour of living creatures is much more complicated; interactions between different individuals and species might be taken into account.

All parts of the definition of the PS presented in the Introduction are equally important.

- i*). The population under consideration might not be very small. Otherwise if calculations are made for the same (small) number of objects, statistical fluctuations will dominate in the final results. If oppositely calculations are made for a large number of objects, then a comparison with a small actual population is not informative. A similar problem can appear even for relatively large populations if a few rare objects can dominate in the final integral value of some parameter. For example a few rare luminous stars (or even one star) can dominate in the integral colour of a cluster (see [6, 7] for a discussion of some limitations of the PS method).

- ii*). An evolutionary path of each object might not be very simple or very complicated. In the first case a PS itself becomes unnecessary, as analytical calculations can be used (however, in early PSs of single and binary stars in the frame of very simplified models authors applied analytical calculations). If an evolutionary track is too complex then it is just difficult to model it, and in addition the number of possible states becomes too large, possibly exceeding the number of objects in the observed population.

- iii*). Objects should not significantly interact with each other, otherwise it is impossible to

follow individual tracks. However, interactions are not absolutely forbidden (for example, stars in globular cluster can interact and form or destroy binaries).

From the philosophical point of view it is possible to divide most of PS calculations into two stages:

- Construction of a model population of objects.
- Calculation of quantities of interest (and their distributions) using the constructed population.

Quantities of interest can include: a number of particular sources, average or extreme values of their parameters and statistical moments of these parameters, correlations between given pairs of parameters, different parameter distributions which can be one-parametric or multi-dimensional, cumulative or in the differential form, etc.

If a set of sources itself (the whole population or its part) is not the main goal of a study, then the set may be constructed implicitly. In such a case a task of calculating of a parameter or its distribution is formally reduced to a multi-dimensional integration of a complicated function which on its part depends on an evolutionary scenario. To calculate such integrals one can use multi-dimensional regular grids or the Monte-Carlo method. Each of this two approaches has advantages and drawbacks.

The way of integration (the classical one or the Monte-Carlo) depends on a method of setting initial parameters of the objects under consideration. A chosen range of initial parameters should cover all the region of initial conditions which is valid for a given problem. Then the coverage should be uniform enough. If an integration proceeds in the classical way, then a chosen region of initial conditions should be covered by a (quasi)rectangular grid, not necessarily uniform. In the Monte-Carlo method initial conditions are chosen stochastically with statistical weights proportional to an assumed actual realization. The second method provides the most realistic populations of modeled objects.

4 Population synthesis of close binary systems

In this section we describe the PS of binary stars. In particular we shall focus on systems with compact objects (neutron stars or black holes). A detailed description of the evolutionary scenario for binary systems can be found in [8].¹ We start with a very brief description of the stellar evolution (details can be found in many standard textbooks, see for example [9] and references therein).

¹This review can be found on the Web at <http://xray.sai.msu.ru/~mystery/articles/review/>.

It is possible to distinguish two main parts of a star's life: when it is a normal non-degenerate star, and when it becomes a compact stellar remnant (a white dwarf, a neutron star or a black hole). Thermonuclear reactions are an energy source of normal stars (see lecture notes by D.K. Nadezhin in this volume). In some sense an evolution of a normal star can be reduced to changes of the nuclear fuel (central hydrogen, hydrogen in a thin layer, helium, etc. up to iron for the most massive stars). About 90% of its life a normal star spends at the main sequence burning hydrogen in its a center. After the main sequence stage a star can significantly change its radius, surface temperature and loose large part of its mass (mass losses at the main sequence are important only for very massive stars). Finally a star becomes a compact remnant. Objects with initial masses $\lesssim 8\text{--}10 M_{\odot}$ become white dwarfs (of course for stars with $M \lesssim 0.8 M_{\odot}$ an evolution prior to a white dwarf formation can be very long, longer than the present age of the Universe). More massive stars explode and form neutron stars or black holes. The boundary which divides neutron star progenitors from black hole progenitors is uncertain. Roughly this value is about $25\text{--}35 M_{\odot}$. On the other hand, there are suggestions that it can depend on rotation and magnetic field of a star (see [10] for a detailed discussion of late stages of the stellar evolution).

The main parameter governing the evolution of a single normal star (and in some sense of a single compact object) is its mass. An isolated star can only loose its mass due to a stellar wind. In a binary the situation is drastically different: a star may become a donor or an acceptor, i.e. it can transfer matter to a second companion or to get mass from it. This option immediately shows that the evolution in a binary is much more complicated.

A small amount of mass can be transferred via a stellar wind, but the main channel of a mass transfer is an accretion due to the Roche lobe overflow. For example, in such a process, a normal massive star can transfer all its outer layers and become a Wolf-Rayet star – a naked helium core. If the secondary star is a compact object, then an accretion results in an X-ray source formation.

Different types of compact objects have distinct evolutionary paths. A black hole can increase its mass due to an accretion in a binary. Because of the same reason it can be also spun-up. Isolated black holes can be called nearly non-evolving objects. For isolated white dwarfs the most important aspect of evolution is cooling. The richest life full of different stages is a fate of neutron stars, especially in binaries.

A neutron star can appear as a visible source due to its thermal emission, due to the radio pulsar activity, due to an accretion onto its surface, or due to an interaction of its magnetic field with a surrounding plasma. A neutron star's surface temperature, its spin period and magnetic field (plus parameters of the surrounding medium) are the main parameters which determine an astrophysical appearance of a neutron star. Therefore it is possible to

distinguish at least two types of evolution of these objects: thermal history and magnetorotational evolution. For some sources these two types of evolution do not influence each other, for some they do (for example for magnetars).

The first PS studies of binary stars were performed at the beginning of 80s [11, 12, 13, 14, 15]. These authors utilized the evolutionary scenario constructed by Paczyński, Kippenhahn, Weigert, Iben and many others (see reviews in [16, 17]).

Here we shall discuss the PS of close binary systems with compact objects (white dwarfs, neutron stars and/or black holes). The evolution of binary stars produces a variety of different objects with different observational manifestations. In this notes we cannot describe all aspects of the problem, so we just shortly mention some types of objects and some observed phenomena. We shall discuss cataclysmic variables, binary white dwarfs, X-ray binaries, millisecond radio pulsars, and ultraluminous sources. We describe results of researches of the following phenomena: a common envelope stage and type Ia supernovae. Then we present main results and difficulties in studies of gravitational wave sources. We discuss coalescence of neutron stars and black holes and gravitation radiation from binaries. In the last part of the section we discuss binary evolution in dense stellar clusters taking into consideration far and close stellar encounters.

Before we start to discuss different types of sources and phenomena let us give a short (and probably incomplete) list of active scientific groups working on the PS of close binaries and references to some of their recent results in order to provide a guide for the readers interested in this subject:

- Portegies Zwart et al.: PS of dense stellar clusters [18, 19, 20, 21], intermediate mass black holes [22, 23], X-ray sources [24, 25], etc.
- Podsiadlowski et al.: evolution of compact binaries and neutron stars kicks [26, 27, 28], hypernovae and gamma-ray bursts [29, 30], other questions [31, 32].
- Kalogera et al.: neutron stars coalescence rate and gravitational radiation [33, 34, 35, 36], radio pulsars, neutron stars and black holes [37, 38].
- Yungelson et al.: double white dwarfs and supernova type Ia [39, 40, 41, 42], stellar gravitational wave sources [43], etc.

4.1 Census and distributions of compact stars

The PS of binaries is a useful tool to study statistical properties of stars including compact objects. We wish to address the following questions:

- How many compact objects of different types are born?
- What is the mass distribution of these compact objects?
- What is the relation between a stellar initial mass and a final mass of a compact object taking into account the binary evolution?
- With a given distribution of compact object masses, what are relative numbers of different types of objects (neutron stars, quark stars, black holes)?

The binary PS is a hot point of astrophysics. Different scientific groups all over the world have their own codes for calculations of the binary evolution, and produce a huge flow of publications on different aspects, discussing all the questions mentioned above (see, for example, [44, 45, 46, 47, 48, 49, 50]). Below we briefly mention some of these results referring to original papers for details.

Pols and Marinus [44] calculate an evolution of young open clusters. The authors focus on blue stragglers. There are several hypothesis on the nature of these sources. They can be binaries consisting of two main sequence star or one main sequence star and a compact object; or they can be single peculiar stars, for example formed by merging of two normal stars. Distributions of age and other parameters are presented.

In [45] the authors consider merging binaries consisting of two neutron stars or black holes. An estimate of a number of binary radio pulsars (systems like the Hulse-Taylor radio pulsar) is made.

Ref. [46] is devoted to formation and evolution of binary neutron stars. The amount of young and recycled binary pulsars is calculated for different assumptions on kicks, initial mass ratio, and presence or absence of hyperaccretion regime.

The main goal of [47] is to study coalescence of binary neutron stars with implications to gamma-ray bursts (now this scenario is mainly applied to short gamma-ray bursts, while long bursts are explained as hypernova events). The authors estimate birth and merging rates, spatial velocities; also they calculate spatial evolution of the binaries and their displacement from the host galaxies.

In [49] the mass distribution of compact objects is calculated for single and binary stars for different types of a secondary component. The authors include into their model a possibility of a quark star formation. However, their results are very model dependent, and in general the mass spectrum of compact objects is poorly known now (see also sec 5.3 below).

At the end of this subsection we want to attract your attention to two important notices. First, simulations considered above use different evolutionary tracks of normal stars (analytical approximations, evolutionary track libraries, etc.), various initial distributions (initial

mass ratio, kicks, magnetic fields, etc.), all sorts of assumptions about complex stages of the evolution (common envelope, accretion induced collapse, merging, etc.). Second, in the final results of different calculations there are some discrepancies, though general predictions are very similar and robust.

4.2 Cataclysmic variables

A cataclysmic variable is a binary system consisting of an accreting white dwarf and a low-mass main sequence star filling its Roche lobe. The state with a Roche lobe overflow is supported by angular momentum losses due to the magnetic stellar wind braking or due to gravitational radiation. Magnetic wind originates due to a presence of a convective envelope of a normal star. This mechanism works for stars in the mass range $0.3\text{--}1.5 M_{\odot}$. When the mass of a donor star becomes smaller than $\sim 0.3 M_{\odot}$ the magnetic braking vanishes and accretion stops. Later the evolution of the binary is determined by the gravitational wave radiation. After a period of an X-ray quiescence the normal star fills its Roche lobe again, and the X-ray source switches on. The following evolution of such object is mainly driven by emission of the gravitational waves.

The magnetic braking accretion is more powerful than accretion driven by gravitational wave momentum losses, and the corresponding sources are brighter. Besides the magnetic braking and gravitational waves there is another way to loose orbital momentum: an interaction with a circumbinary disc. Ordinarily this mechanism is less effective.

The model described above predicts a gap in the orbital period distribution. This is really observed at $P_{\text{orb}} \sim 3$ hours. Almost all long period cataclysmic variables are induced by the magnetic stellar wind, all short period ones ($P_{\text{orb}} < 3$ hour) are gravitational wave induced.

A PS study of cataclysmic variables and their orbital period distribution in a classical approach was performed in many publications (see for example [51, 52, 53]). The authors are able to reproduce the observed orbital distribution of the variables. The PS of cataclysmic variables, surrounded by a circumbinary disc is performed in [54, 55]. The authors come to the conclusion that the bimodal period distribution in the frame of the outer disc model is possible only if a proper bimodality of disc feed properties is presented.

4.3 Binary white dwarfs

In applications of a PS to binary white dwarfs there are two topics which seems to us the most interesting.

At first, the amount of very close binary white dwarf systems can help to constrain the efficiency of angular momentum losses at the common envelope stage. Most of the close

binary white dwarf systems are formed after two common envelope episodes. The first one is due to the Roche lobe filling by the primary, and the second starts after the secondary component also fills its Roche lobe. This type of systems is the most sensitive indicator of common envelope parameters. According to [56, 57] the parameter of the efficiency is $\alpha_{\text{CE}} \simeq 0.3\text{--}0.5$. This value does not coincide with results of other PSs calculations which take into account evolution with just a single common envelope stage [58].

Coalescence of binary white dwarfs is the second important topic. This process is one of evolutionary channels leading to a type Ia supernova explosion – so-called double-degenerate scenario.² This topic was studied, for example, in [33, 34, 59, 36, 42, 35]. It is shown that coalescence of white dwarfs is able to produce type Ia supernovae with a rate $\sim 10^{-3} \text{ yr}^{-1}$. This estimate is valid for both old and young stellar populations (i.e. for elliptical and spiral galaxies). For the second channel (accretion onto a white dwarf from a normal companion) calculations give an estimate of the supernova Ia rate $\sim 10^{-4} \text{ yr}^{-1}$. However, it is worth noting that recent studies show that the supernova Ia rate per unit mass is much higher for young stellar populations [60]. This can be an argument against the double-degenerate scenario.

4.4 Accreting X-ray sources and millisecond pulsars

Present day orbital X-ray observatories like Chandra and XMM have a sufficient sensitivity to detect all bright binary X-ray sources in the Galaxy and almost all sources with moderate luminosities. There are many different types of binary X-ray sources with permanent or transient hard emission: X-ray pulsars, X-ray bursters, so-called “atoll” and Z-sources, sources with QPO, X-ray novae, black hole candidates etc. Some sources may display several of the enumerated phenomena simultaneously. Observational properties of different types of the sources are discussed in the following reviews [61, 62, 63, 64, 65, 66].

The first PS investigations of high-mass X-ray binaries were made about 20 years ago. PS studies of low and intermediate-mass X-ray binaries with more complex evolution started later. Recently different authors started to include into calculations individual properties of studied regions; for example, starformation history in a given galaxy, or mass and metallicity of a given region of starformation. Below we give a short list of recent publications devoted to PS of binary X-ray sources.

In the papers [67, 68, 69] and in a series of papers [69] the authors calculate accretion rates, luminosities, spatial velocities, and birth rates of massive X-ray binary sources with supergiants and Wolf-Rayet companions. In [68] a population of X-ray sources with Be-

²The second possibility to produce such supernova is the accretion induced collapse of a white dwarf – the single-degenerate scenario.

companions is studied. In [67] – a population of the Thorne-Zytkow objects (red giants with neutron star core – a possible result of merging of components of a massive binary during a common envelope stage). In [70] a population of X-ray sources near the center of our Galaxy is synthesized. Comparison of observed luminosity distributions of X-ray sources with PS results of [71] indicates the presence of several populations of X-ray binaries in nearby galaxies. Each population has its own formation and evolutionary history, depending on the host environment.

In [72] the authors try to reproduce about 140 observed low-mass binary galactic X-ray sources in the frame of low and intermediate-mass binary evolutionary scenario. Ref. [73] represents a more advanced version of the previous PS by the same team. In this publication properties of millisecond pulsars are also considered.

In the end of this subsection we want to emphasize two interesting, in our opinion, groups of X-ray sources.

Various X-ray systems are different not only in a physical sense, but also in PS contents. Equal observational amounts of sources may be produced by numerous short lived objects or by rare long lived ones. A PS of the latter objects is a sufficiently more difficult task. The most interesting representatives of that sort of objects are accreting progenitors of millisecond radio pulsars. A PS of millisecond pulsars and their progenitors is discussed in [74, 75].

X-ray novae are accreting close binaries consisting of a black hole and a low mass main sequence component. To create such a system it is necessary to start with a binary with very high initial mass ratio. A black hole progenitor have to be at least as massive as $25\text{--}30 M_{\odot}$. Of course such a massive star loses significant part of its mass prior to a compact object formation. A low-mass component saves its initial small mass. It is not easy to reproduce a significant number of X-novae in the frame of a standard scenario of binary evolution. In 1986 Eggleton and Verbunt [76] suggested a scenario for creation of an X-ray nova from a massive hierarchical triple system. The evolutionary scenario for hierarchical triple and multiple stellar systems posses sufficiently more potential variety instead of the binary evolution scenario (see, for example, [77]). Computer realization of an evolution of a triple system is very complex, At present we know only one realization of such code. Comparison of these two scenarios for the X-ray nova evolution and results of binary and triple PS calculations can be found in [78].

4.5 Ultraluminous X-Ray sources

Ultraluminous X-ray sources are very bright point X-ray objects outside nuclear regions of host galaxies (see a catalogue in [79] and a recent review in [80]). Ordinarily this objects are defined as sources with an X-ray luminosity larger than $10^{39} \text{ erg s}^{-1}$ (from $\sim 10^{39}$ up to

$\sim 10^{42} \text{ erg s}^{-1}$). They were discovered by Einstein space observatory [81]. Large number of these sources was later found by ROSAT and Chandra. In our Galaxy bright ultraluminous sources are absent.

There are two popular models of ultraluminous sources. In one of them an accretion with nearly Eddington rate proceeds onto an intermediate mass black hole ($10^2\text{--}10^3 M_\odot$) in a binary system (e.g. [82], see [83] for a detailed discussion on the properties of intermediate mass black holes). In another hypothesis it is supposed that we deal with normal stellar mass black holes, but radiation is not symmetric, being beamed into two jets. If we look along the jet, then we observe a very bright source. In this model ultraluminous sources may be just a high luminosity tail of the stellar-mass black hole luminosity distribution (e.g. [84, 85]). Both hypothesis have advantages and drawbacks. Probably the ultraluminous sources population is formed by two (or even more?) different types of objects [86].

Podsiadlowski et al. [87] perform a systematic study of formation and evolution of black hole binaries and find that indeed their models are consistent with the observed luminosity function, and a typical number of ultraluminous sources per galaxy can be naturally explained. Note, that as the ultraluminous sources are very rare, bright sources of that type are observed with a rate 0.01 per galaxy, calculations of their luminosity function and its comparison with observations is not trivial. Different statistical and selection effects might be taken into account.

4.6 Stellar gravitational wave background

New generations of ground-based (LIGO, VIRGO, GEO) and space (LISA) gravitational wave interferometric detectors will be sensitive at $10^1\text{--}10^3$ and at $10^{-4}\text{--}10^{-1}$ Hz, respectively. In this frequency bands there are three general types of stellar gravitational wave sources ³:

1. Core-collapsed supernovae (type II, Ib or Ic);
2. Rapidly rotating non-axisymmetric neutron stars;
3. Close binaries and coalescing compact stars.

Let us briefly discuss these items (coalescence of compact objects is also a subject of the next subsection). The rate of supernovae is roughly known ($\sim 1/30\text{--}1/50 \text{ yrs}^{-1}$). Nevertheless, to calculate a possible rate of detection of gravitational signals it is necessary to know how much energy is transferred into gravitational radiation during an explosion. As a complete self-consistent theory of supernovae is absent, the fraction of the energy of an explosion

³We do not discuss coalescence of supermassive black holes, which can be an important type of sources for LISA.

which goes into gravitational waves is unknown. This is the most important uncertainty of this subject.

Rapidly rotating neutron stars emit monochromatic gravitational waves. The energy flux may depend on a proper non-sphericity of a neutron star. For very fast neutron stars the symmetry breaking appears due to the Chandrasekhar-Friedman-Schutz instability [88, 89], or due to an r-mode instability [90, 91]. In this case the flux depends on viscosity and temperature of a neutron star crust and interior. All parameters here, especially viscosity, are not well known. Results of different studies are still very controversial.

The gravitational radiation from binary stars is well understood and inevitable in the frame of the General Relativity. The first estimate of the gravitational wave background from galactic binaries was obtained by Mironovskij in 1965 [92] on the base of counts of W UMa systems. This author got an estimate of the dimensionless amplitude of metric perturbations equal to $h \sim 10^{-21}$. The spectrum of the stellar galactic gravitational wave background was calculated with the binary PS model in 1987 [93]. The first calculations show a presence of two different parts in the spectrum. The first one, at low and intermediate frequencies with a maximum at $\nu \sim 10^{-4}$ – 10^{-3} Hz, is linked with the gravitational radiation of binaries. The second represents a high frequency tail of the spectrum. It appears due to merging of white dwarfs and neutron stars. Recent results on this subject can be found in [94, 95, 96].

4.7 Merging and coalescence rate of neutron stars and black holes

Components of a relativistic binary system with an orbital period less than ~ 14 hours can spiral-in and merge in the Hubble time due to the gravitation wave emission. Coalescence of neutron stars was widely discussed as a possible source of gamma-ray bursts and as a source of strong gravitational wave pulses detectable by ground-based detectors.

The rate of neutron stars coalescence is a key parameter in modeling of an observable rate of gravitational wave bursts. There are two different estimates of this value: one is based on PS calculations, and another is based on the number of observed binary radio pulsars with short orbital periods (like the famous Hulse-Taylor system). During last twenty years there was some contradiction between these two approaches. Since the first paper [93] and later on (see [97] and references therein) PS models predict the coalescence rate about 10^{-4} yr^{-1} for a spiral galaxy similar to the Milky Way. The direct estimate based on the binary radio pulsar statistics predict significantly lower coalescence rate $\sim 10^{-6} \text{ yr}^{-1}$ [98]. Later this value has been increased up to $2 \cdot 10^{-5} \text{ yr}^{-1}$ (for example [99, 35]), nevertheless, the difference did not disappear. This method takes into account only binaries where at least one component is observed as a radio pulsar. In a PS model all merging neutron stars are

taken into account, including radioquiet ones. This difference can partially explain the lack of convergence between the two estimates. The discovery of the double binary radio pulsar J0737-3039 with a very short orbital period [100] increases the observational prediction by nearly an order on magnitude, and so now the problem is almost solved [101]. The history of the question is briefly described in [102]. One can follow the references given in that paper.

If one wants to predict a detection rate of gravitational wave signals from coalescing compact objects it is not enough to know the rate of this events. It is also necessary to take into account the power distribution of the sources, i.e. their luminosities in gravitational waves. For a fixed sensitivity of a gravitational wave antenna the maximum distance from which we still can register a coalescence depends first of all on a mass of a merging binary (more precisely on the so called “chirp mass” – some combination of masses of the binary components). If ground-based interferometers can reach their nominal sensitivity they can detect neutron star merging from a few dozen megaparsec. The coalescence rate of binary black holes is probably lower by 1-2 order of magnitude in comparison with neutron star systems, but note that black holes are more massive (at least by a factor a few). Due to strong dependence of the maximum detection distance on a binary’s mass we can detect an event from sufficiently larger volume of space. This factor can be even more important and the *detection rate* of black holes coalescence can exceed the same quantity for binary neutron stars [8]. I.e. the first detected signals can be due to coalescence of a black hole with a neutron star or with another black hole!

Note, that values of coalescence rates are very sensitive to details of evolutionary scenario. For example, as progenitors of black holes are very massive stars they should form relatively wide binaries. If an orbit of a system is not changed during a black hole formation then the time prior the components merge is very long. Thus in this case the detection rate is zero. To change the situation it is necessary to introduce a natal kick for black holes [103]. These is quite possible if a black hole is formed in a two-stage collapse. Recent studies of black hole binaries support the non-zero value of the kick velocity [104].

4.8 Binaries in dense stellar clusters

Evolution of binary populations in dense stellar clusters (globular clusters and central, i.e. circumnuclear, clusters in galaxies) differs from evolution of the same binaries in the galactic field. There are two reasons for the difference: an interaction of individual binaries with gravitational field of a cluster and a close fly-by (a close passages of a star near a binary system).

The first effect makes close binaries (so called “hard” systems) closer, while wide systems become wider. The boundary between the two types of systems is determined by the

condition $v_{\text{orb}} \simeq v_{\text{cluster}}$. Here v_{orb} – orbital velocity of a binary, v_{cluster} – mean velocity of stars in a cluster.

There are various effects of a close encounters. A star flying-by can replace one of the binary component, can disrupt the binary, can induce a merging of the binary, or can just change binary orbital parameters. Probabilities of all of these effects were studied in [105, 106]. If a binary interact with another binary system, then, of course, there are more possible outputs of the interaction (see [107, 108]).

Taking into account the possibility of a close fly-by, i.e pair stellar interaction, we destroy one of the basic PS principle about an independence of objects' evolution. However, if point-to-point interaction remain rare, we still can use a PS in some modified form.

There are two possible ways to modify the PS scheme to allow for close encounters. One method is to use a complete binary evolution scenario with simplified stellar dynamics. Another one, vice versa, includes precise N-body calculations of stellar motions with a reduced evolutionary description. The second version was realized in a recent series of papers by Portegies Zwart, Aarseth et al. [109, 110, 19, 111, 108]. Although this is just a first step in realization of a full scenario of evolution in dense clusters.

5 Population synthesis of isolated neutron stars

In this section we discuss a PS of isolated neutron stars (NSs).⁴ We focus on three types of research: radio pulsars studies, modeling of old NS evolution, and studies of close-by cooling NSs. The first two include as the main ingredients initial period, velocity, and magnetic field distributions together with evolutionary laws for all these quantities. In addition such a parameter as the angle between spin and magnetic axes can be included. For the PS of cooling NSs ingredients are quite different, as the thermal history is not tightly connected with periods and magnetic fields, instead the mass spectrum of NSs is important.

5.1 Radiopulsars

As radio pulsars are the most known example of NSs and form the largest observed population of these objects, it is not a big surprise that there were several PS studies dedicated to these sources. The crucial feature of such models is the necessity of simulating of the detectability of radio pulsars, i.e. it is not enough to calculate an artificial population of NSs, it is also necessary to imitate a radio survey with realistic parameters to compare calculations with observations.

⁴Russian-speaking readers can find more information on this subject in our review [112]. It is also available on the Web at <http://xray.sai.msu.ru/~polar/html/kniga.html>.

It is possible to distinguish three main reasons for a PS of radio pulsars. The first one is related to a determination of initial parameters of NSs (periods, magnetic fields, velocities). The second deals with parameters of the evolution (for example the magnetic field decay). And finally the third task is to predict properties of some unobserved population (highly magnetized pulsars, gamma-ray pulsars, etc.) basing on some known initial and evolutionary parameters.

One of the key questions concerning the evolution of the population of radio pulsars is the following: *does the magnetic field of NSs decay?* Bhattacharya, Hartman, Verbunt and Wijers address this question in a series of papers (see [5] and references to earlier papers therein). They use Monte-Carlo simulations to reproduce distributions of radio pulsars on the $P - \dot{P}$ plane. Also the authors calculate velocity distribution of NSs, their spatial distributions in different projections, and dispersion measure distributions. Verbunt et al. assume initial magnetic field distribution of NSs similar to the one observed for radio pulsars: a gaussian in a logarithmic scale. They also take into account observability of the modeled sources. These authors conclude that the best fit includes the magnetic field decay on the time scale $\sim 10^8$ yrs, and that significant amount of NSs are born with velocities below 200 km s^{-1} .

There are several hypothesis about initial conditions and evolutionary laws made by these authors that can be criticized. Both velocity distributions used by the authors are single-mode ones. Nowadays the bimodality of the velocity distribution is considered to be a standard assumption. Also modern observations strongly suggest that the number of high velocity NSs is larger than it follows from the distributions used by Verbunt et al. The initial magnetic field distribution used by the authors hardly can explain the existence of magnetars. Finally Verbunt et al. do not discuss the rôle of the inclination angle χ (between spin and magnetic axes) and its evolution.

A more updated PS of radio pulsars is performed by Regimbau and de Freitas Pacheco [113]. They include into the model the evolution of χ and discuss the appearance of magnetars. A comparison with the papers by Bhattacharya et al. shows how changes in parameters of a model of the PS change the final results. Regimbau and de Freitas Pacheco obtain an agreement of the observed data with a model without the magnetic field decay, but with an inclusion of an evolving (increasing) angle χ . Another important conclusion reached in [113] is related to magnetars. The authors obtain that about 23% of NSs are highly magnetized ($B > 10^{14} \text{ G}$). Their results suggest that there is no bimodality in the magnetic field distribution, magnetars naturally appear from the high field tail of the distribution which also includes radio pulsars.

A separate topic of radio pulsar studies is related to γ -ray observations. Gonthier et al.

[114] address the question of active radio pulsar detection in the high-energy band. Such a research is very relevant now in the light of future space projects as AGILE and GLAST. In the study by Gonthier et al. several new important parameters appear as the authors incorporate independent models for the radio and gamma-ray beam geometry.

An unexpected result obtained by these authors is the necessity of the magnetic field decay with relatively short time scale – 2.8 Myr. Predictions for AGILE and GLAST observations are very optimistic. Gonthier et al. find that GLAST can observe up to 600 NSs in gamma-rays.

Beskin and Eliseeva [115] study a possibility of observation of dead pulsars as γ -ray sources. Strictly speaking these authors do not perform a PS study, nevertheless their work can be considered as an important input for population models. They obtain a period distribution of dead (i.e. non-active) pulsars for two models of the evolution of the angle χ between spin and magnetic axes. Up to our knowledge a PS of NSs which are not active radio pulsars with a detailed model of the evolution of χ has never been performed. Such study can be very important as it can change conclusions about the stage distribution of old NSs which we are going to discuss in the next subsection.

5.2 Neutron star census and isolated accretors

Since early 70s different authors discuss a possibility of observations of isolated accreting NSs. During these years conclusions changed from very optimistic (nearly all old NSs are accreting; it is possible to observe thousands of them with X-ray satellites like ROSAT) to very pessimistic (only a small fraction of isolated NSs can reach the stage of accretion; most of accretors are very dim sources). In 90s Monte-Carlo simulations of the NS population were performed by several groups. We mention here two illustrative papers by Blaes and Rajagopal [116] and Manning et al. [117].

The authors of [116] are motivated by a discussion on the nature of gamma-ray bursts. Their PS study has a goal to calculate the distribution of masses accreted onto isolated NSs in the solar neighborhood. The main ingredients of the model are: initial spatial distribution of NSs, their initial velocity distribution, the galactic potential, interstellar medium (ISM) distribution, type of accretion. Blaes and Rajagopal (as many other researchers before mid-90s) assume that all NSs relatively quickly start to accrete. This assumption is the main mistake of many models at that period, as the evolution of NSs has been absolutely ignored. Due to this approach all groups obtained a huge number of accreting isolated NSs.

We want to make one very important comment concerning the evolution of isolated NSs. Before 1994 it was assumed that the velocity distribution of NSs allows an existence of a big number of low-velocity compact objects. After the paper by Lyne and Lorimer [118]

has appeared it became clear that quite oppositely NSs (at least radio pulsars) are very high-velocity objects. The present day velocity distribution [119] is bimodal with more than 1/2 of NSs with velocities larger than $500\text{--}600 \text{ km s}^{-1}$. Very often different authors claim that small (actually zero) number of discovered isolated accretors can be explained by low luminosities due to high velocity of NSs (in the Bondi formula the accretion rate is inversely proportional to a cube of velocity). In some sense this statement can be put in such a way: there are a lot of accretors, but with modern satellites we cannot detect them. However, it is not the case! NSs with big initial velocities just never reach the stage of accretion. The number of isolated accretors is as low as the number of low velocity NSs.

Manning et al. [117] discuss the duration of stages preceding the onset of accretion. On the other hand, they do not use the new (Lyne and Lorimer) velocity distribution. That is why they find that a significant fraction of NSs could reach the accretion stage in a lifetime of the Galaxy. Anyway, as Manning et al. included the evolution before accretion starts, probably their analysis was the most complete till the end of 90s.

An attempt to model a global evolutionary distribution of isolated NSs is made in [120]. This *neutron star census* is performed to obtain fractions of NS population on each of the four main evolutionary stages: *ejectors*, *propellers*, *accretors*, *georotators* (see [8] for the stages description). From the present day point of view this study has several drawbacks. The authors use a very simple initial period and magnetic field distributions. The velocity distribution is just a single-mode one. The propeller and georotator stages are treated in a simplified way. The rôle of χ is not studied. Nevertheless, in our opinion, the main features of the realistic distribution are picked up correctly. For standard magnetic fields and without decay a NS spends most of its life as an ejector (or as a georotator if velocity is high). For decaying fields the propeller stage can be the most populated.

In [120] it is found that for reasonable velocities more than 90% of isolated NSs spend all their lives as ejectors. Fractions of propellers and georotators are found to be small. However, we note that the authors treat the georotator stage in a simplified manner, and depending on definitions it is possible to say that significant fraction of very high-velocity NSs are georotators. Also the stage of subsonic propeller (see [121] and references therein) is not taken into account.

A more detailed study of properties of accreting isolated NSs is performed in [122]. The authors obtain the $\text{Log } N - \text{Log } S$ distribution for accretors. It is shown that at low fluxes (say, below $10^{-13}\text{--}10^{-14} \text{ erg s}^{-1}$) accretors can become more abundant than coolers (see the next subsection on that type of sources). Also velocity, temperature and accretion rate distributions of isolated accreting NSs are derived.

To conclude this subsection we want to stress once again that PS models for old isolated

NSs passed a long way. Many parameters were significantly refined. Nevertheless, as details of magneto-rotational evolution of NSs are still unknown (evolution of χ , properties of the propeller stage, accretion rate and efficiency, etc.) in future more detailed calculations will inevitably appear.

5.3 Close-by young neutron stars

Quite a different branch of NS population studies is related to young close-by cooling objects – *coolers*. Presently, thanks to the ROSAT satellite, we know a set of nearly a dozen young hot close-by NSs of different nature (see the list in [123]). Among these sources there are four normal radio pulsars, Geminga and a geminga-like object, and seven radioquiet NSs (so-called "Magnificent seven"). Of course, this set of sources is not complete: there should be more objects of that kind in the solar vicinity, it is just very difficult to identify them in the crowded environment close to the Galactic plane. Nevertheless, as all the sources were detected by one instrument (and if possible we select them with upper bounds on age and distance from the Sun) this set is uniform enough to perform a PS not worrying a lot about selection effects (but see below on possible correlations between different parameters of NSs). The goal of such a study can be twofold: to understand the origin of an observed population of close-by *coolers*, and to test the theoretical cooling curves by the observed objects. The former approach was used in [123, 124]. The latter one – in [125].

The main ingredients of such studies include:

- Initial spatial distribution of NSs (or, say, of progenitor stars);
- Initial velocity distribution of NSs;
- Mass spectrum of NSs;
- Cooling curves;
- Properties of NS emission;
- Properties of the ISM (to take absorption into account).

Some of the ingredients are known well enough for the purposes of this particular problem (spatial and velocity distributions, and the ISM properties). Others are poorly known.

In [123] it was shown that the local population of hot NSs from which the thermal emission is detected is genetically connected with the Gould Belt. This structure is responsible for the overabundance of massive stars (and so of young compact objects) in the Solar vicinity.

The authors of [125] suggested to use $\text{Log } N - \text{Log } S$ distribution of close-by coolers as the second additional test for calculations of the NS thermal evolution. Indeed, if all but one ingredients of the PS are known, then comparison with an actual data can be used to test the remaining ingredient.

Presently the only way to compare calculations of the thermal evolution of NSs with observational data is the so-called T-t (temperature vs. age) test. Being the most natural test for this problem it has some drawbacks:

- There are uncertainties in temperature and ages of observed sources;
- The test is not sensitive enough for ages $> \text{few } 10^5$ yrs;
- The set of objects is not uniform at all.

$\text{Log } N - \text{Log } S$ as the second test has a complimentary to the T-t test set of advantages and disadvantages.

- There is no uncertainty in observational data;
- The set of sources is nearly uniform;
- The test is mainly sensitive to properties of NSs with ages $> \text{few } 10^5$ yrs.

On the other hand, there are some drawbacks (see [125] for more details).

Preliminary results show that the $\text{Log } N - \text{Log } S$ test works well as an addition to the T-t test (we note, that the idea is to use the two tests together, not $\text{Log } N - \text{Log } S$ alone). This is a good example how an astrophysical PS study can be used to test predictions of physics of matter in extreme states.

5.4 Unsolved problems

Many ingredients of the PS models described above are not well known.

In our opinion the main difficulty of the radio pulsar PS is connected with the fact that we do not understand well enough emission mechanisms in different energy ranges. Due to this it is difficult to derive many crucial parameters and to produce results which can be directly compared with observations. Many authors obtain drastically different results even if all parameters of a model are the same, but mechanisms of radio pulsar emission are different. As a rule such parameters as initial period and magnetic field distributions are not well known. Evolutions of the magnetic field and of the angle between spin and magnetic axes are unknown too.

A PS of old NSs suffers from the lack of observational data. No old isolated NS which is not a millisecond radio pulsar is known, and there is very little hope to find a way to observe an isolated NS at a stage of propeller or georotator. Among theoretical uncertainties in addition to the ones described in the previous paragraph it is necessary to mention difficulties with the description of the propeller stage. Also the regime of accretion onto an isolated NS and its efficiency is not known [126], estimates made with the Bondi formula can be just an upper limit.

Calculation of the number of observable coolers is the subject of uncertainties in the mass spectrum of NSs, in cooling curves and in emission properties (spectrum and surface emissivity).

6 Other examples of population synthesis

6.1 Stellar populations and spectral studies

The story of PS of stellar populations started in early 70s (see for example [4] for early references and for a brief historical introduction). Historically the *empirical population synthesis* appeared earlier. Nowadays both methods are actively used. To learn more about the *evolutionary population synthesis* we can recommend the paper [4] and references therein. The *empirical population synthesis* is well described in [2]. Both methods together are discussed in [1].

Both methods are mainly motivated by the same problem: to derive properties of a stellar content (ages, chemical composition, etc) basing on an integral spectrum of a galaxy (an inverse problem, to predict a spectrum of a galaxy, is also possible). The *empirical population synthesis* is based on a *decomposition* of a galactic spectrum to known spectra of stars of star populations. I.e. it is necessary to find a linear combination of known spectra which fits an observed one (very often equivalent widths of spectral lines are used as quantities to fit). The main problem of the method is connected with non-uniqueness of a solution due to a number of free parameters. For example, there is a known degeneracy between age and metallicity: changes in metallicity can mimic a change in the age of a population. An increased metallicity and an increased age both make the integrated spectral energy distribution redder (see for example a discussion in [127]).

The main ingredients of the *evolutionary PS* are:

- Initial mass function;
- Starformation rate;
- Initial chemical composition and a rate of chemical enrichment.

Main problems are connected with non-sufficiently understood stages of the stellar evolution – stages after the main sequence (see for example [3]). Also there is a degeneracy between age and initial mass function (it can be important for age determinations of starbursts [128]), for example, a smaller fraction of massive stars can mimic an effect of aging of a stellar population.

Both methods use some input spectral information. In the case of the *evolutionary PS* it includes: spectra parametrized by chemical composition, surface gravity and effective temperature. For the *empirical PS* input data contains spectra of some types of stellar populations, for example, spectra of clusters of stars (so-called *single stellar populations*) with the same chemical composition. Ages of stars belonging to each cluster are the same, but ages of different clusters are different, so one can fit an observed spectrum by a composition of star clusters born at different time.

For years the PS of normal stars was done without taking into account the rôle of binary stars. Only in late 90s several groups of authors started to include binaries in their models (see [129] and references therein). Among recent papers we want to mention [130] and a review in [131].

All the authors who studied the influence of binary systems on the integral spectral characteristics in the visual waveband conclude that its rôle is very important. Especially it is important if Wolf-Rayet stars can form only in binaries (an alternative view requires too strong stellar winds for single stars).

Another specific problem is connected with galaxies with high rate of star formation [132]. Modeling young starbursts is a knotty problem due to several reasons:

- Spectra are very complicated as still there are many bright massive stars.
- As stars were not born simultaneously stellar ages are uncertain and the level of this uncertainty is about stars ages.
- There is a lot of dust and the interstellar medium is very inhomogeneous.

An age determination for young star forming systems is discussed in some details in [128]. The approach is based on single stellar populations, with addition of different (depending on time since beginning of a starburst) age indicators. Depending on an age of a population different spectral lines (or other features) can be used as such indicators. For example, for very young stars (~ 4 Myr) nebular emission lines and UV stellar-wind lines can be used as age discriminants.

An example of determination of an age of an old system and references for early papers are given in [127]. These authors also develop a model based on single stellar populations

for ages 1-19 Gyr. Applying the model to the galaxy M32 (a satellite of the Andromeda galaxy) they obtain that it can be described as a single population with a solar metallicity and an age ~ 6.5 Gyr.

Evolutionary synthesis was used in [133] to obtain such an important characteristic as mass to light (M/L) ratio for galaxies. This quantity is widely used to estimate a mass of a galaxy knowing its luminosity in some band.

6.2 Active galactic nuclei and X-ray background

Observations aboard ROSAT and, especially, Chandra satellites have shown that more than 60% (up to 90% in softer X-rays) of the X-ray background (XRB) could be resolved, i.e. point sources are responsible for that radiation. According to optical identifications most of these objects are active galactic nuclei (AGNs) with small addition of clusters of galaxies and stars. Such observations rised interest to PS studies of AGNs.

Initially PS studies of the XRB were motivated by the so-called unified model of AGN (see [134, 135] for examples of PS calculations and for references to earlier papers). In this model all AGN activity is explained by accretion onto central supermassive black hole (see [136] for the unified model description). Different types of AGNs appear due to different black hole masses, different properties of the accreting medium and, especially, due to obscuration by a torus surrounding the black hole. Later on PS models became more phenomenological as all main ingredients are taken from observational data (for example [137, 138, 139]).

Among the ingredients we can mention:

- Relative fraction of obscured and non-obscured AGNs;
- Luminosity functions of AGNs;
- Spectral energy distribution;
- Evolution of these parameters (if any).

Usually models are closer to those called “population synthesis” by Fritze - v. Alvensleben [1] than to the scenarios described in sec. 4–5 (which must be called “evolutionary synthesis” in her notation). In studies of the XRB authors do not follow the evolution of AGNs starting with some initial conditions (seed black holes in centers of galaxies, or even from the scenario of hierarchical clustering). In most of papers the XRB is explained by contribution of absorbed (type 1) and non-absorbed (type 2) AGNs. Spectra of type 2 AGNs can be calculated from spectra of the type 1 by adding some obscuration. This is the main contribution of the unified model to the PS scenario. In early papers the ratio of numbers

of type1 to type2 sources was also taken as the unified model prediction, but now it is taken from the observational data.

Nowadays these models can more or less successfully explain the main properties of the XRB. Nevertheless, it is necessary to make them less phenomenological by linking the PS calculations with calculations of the early evolution of the galaxies (including galaxy formation, seed black holes, etc.). Also the rôle of starburst activity in the XRB is not known well enough (see [139]).

7 Conclusions

In this lecture notes we briefly described the main applications of *population synthesis* in astrophysics. The spectrum of applications is very wide: from normal stars to compact remnants, from integral spectra of galaxies to X-ray background. We mainly focused on evolution of isolated and binary compact objects as these subjects are closer to our personal scientific interests.

The population synthesis as a method of theoretical (numerical) studies is a typically astrophysical tool as in astronomy very often we can observe just a tiny fraction of a population of sources, or cannot resolve single sources at all. In these cases the population synthesis can help to derive or check parameters of the whole set of studied objects, or to predict properties of an unobserved fraction of the sources.

As our knowledge about some population of objects grows models of the population synthesis become more complicated. Finally, when the whole population can be studied observationally, and also an evolution of its members can be directly derived from actual data, population modeling becomes unnecessary (except to show self-consistency of results of observations). However, in most of astrophysical situations we are very far from this ideal, and so the population synthesis calculations will be an important tool for a long time.

Acknowledgments

We want to thank the Organizers of the School for financial support and hospitality. S.P. acknowledges a postdoctoral fellowship from the University of Padova where he worked at the time of the School. This work was partly supported by RFBR grants 04-02-16720 and 03-02-16068.

References

- [1] U. Fritze-v.Alvensleben, in *Stars, Gas and Dust in Galaxies: Exploring the Links*, ASP Conference Proceedings, **221**. Edited by Danielle Alloin, Knut Olsen, and Gaspar Galaz, p.179, e-lanl: astro-ph/0009290 (2000).
- [2] R. Cid Fernandes, L. Sodré Jr., H.R. Schmitt, and J.R.S. Leão, MNRAS **325**, 60 (2001).
- [3] C. Maraston, MNRAS in press, e-lanl: astro-ph/0410207.
- [4] G. Bruzual, and S. Charlot, MNRAS **344**, 1000 (2003).
- [5] F. Verbunt, J.W. Hartman, D. Bhattacharya, R.A.M.J. Wijers, and G. Nelemans, in *Pulsar timing, general relativity and the internal structure of neutron stars*, Edited by Z. Arzoumanian, F. Van der Hooft, E. P. J. van den Heuvel, p.215 (1999).
- [6] M. Cerviño, and D. Valls-Gabaud, MNRAS **338**, 481 (2003).
- [7] M. Cerviño, and V. Luridiana, A&A **413**, 145 (2004).
- [8] V.M. Lipunov, K.A. Postnov, and M.E. Prokhorov, Astrophys. and Space Science Rev., **9**, 1 (1996).
- [9] G.S. Bisnovatyi-Kogan, “Stellar physics”. Berlin: Springer, Astronomy and astrophysics library (2002).
- [10] S.E. Woosley, A. Heger, and T.A. Weaver, Rev. Mod. Phys. **74**, 1015 (2002).
- [11] V.G. Kornilov, and V.M. Lipunov, Soviet. Astron., **27**, 163 (1983);
V.G. Kornilov, and V.M. Lipunov, Soviet. Astron., **27**, 334 (1983).
- [12] V.M. Lipunov, and K.A. Postnov, Sov. Astron. **31**, 288 (1987);
V.M. Lipunov, and K.A. Postnov, Sov. Astron. **31**, 404 (1987).
- [13] R.J. Dewey, and J.M. Cordes, ApJ **321**, 780 (1987).
- [14] M. Manteiga, R.C. Martinez, and A.J. Pickles, Ap&SS **156**, 169 (1989).
- [15] L.R. Yungelson, and A.V. Tutukov, in *Wolf-Rayet Stars and Interpretation of Other Massive Stars in galaxies*, Edited by K.A.Van, B.Hidayet, Dordrecht, Reidel, p.459 (1991).
- [16] B. Paczyński, ARA&A **9**, 183 (1971).

- [17] E.P.J. van den Heuvel, in *Neutron Stars: Theory and Observation*, Edited by J.Ventura, D.Pines, Kluwer Academic Publishers, Dordrecht, p.171 (1991).
- [18] S.F. Portegies Zwart, J. Makino, S.L.W. McMillan, and P. Hut, *ApJ* **546**, L101 (2001).
- [19] D. Pooley, W.H.G. Lewin, and S.F. Anderson, *ApJ* **591**, L131 (2003).
- [20] A. Sills, S. Deiters, P. Eggleton, et al., *New Astron.* **8**, 605 (2003).
- [21] S.F. Portegies Zwart, P. Hut, S.L.W. McMillan, and J. Makino, *MNRAS* **351**, 473 (2004).
- [22] S.F. Portegies Zwart, J. Dewi, and T. Maccarone, *MNRAS* **481** (2004) in press; e-lanl: astro-ph/0406550.
- [23] S.F. Portegies Zwart, e-lanl: astro-ph/0406550.
- [24] G.E. Brown, C.-H. Lee, S.F. Portegies Zwart, and H.A. Bethe, *ApJ* **547**, 345 (2001).
- [25] G. Nelemans, L.R. Yungelson, and S.F. Portegies Zwart, *MNRAS* **349**, 181 (2004).
- [26] E. Pfahl, S. Rappaport, Ph. Podsiadlowski, and H. Spruit, *AAS* **199**, 9508 (2001).
- [27] E. Pfahl, S. Rappaport, and Ph. Podsiadlowski, *ApJ* **571**, L37 (2002).
- [28] P. Podsiadlowski, et al., *ApJ* **612**, 1044 (2004).
- [29] P. Podsiadlowski, M.J. Rees, and M. Ruderman, *MNRAS* **273**, 755 (1995).
- [30] P. Podsiadlowski, et al., *ApJ* **607**, L17 (2004).
- [31] M.E. Beer, and Ph. Podsiadlowski, *MNRAS* **335**, 358 (2002).
- [32] Z. Han, and Ph. Podsiadlowski, *MNRAS* **350**, 1301 (2004).
- [33] V. Kalogera, and D.R. Lorimer, *ApJ* **541**, 319 (2000).
- [34] V. Kalogera, and K. Belczynski, *AIPC* **575**, 107 (2001).
- [35] C. Kim, V. Kalogera, and D.R. Lorimer, *ApJ* **584**, 985 (2003).
- [36] V. Kalogera, et al., *ApJ* **601**, L179 (2004).
- [37] C.L. Fryer, and V. Kalogera, *ApJ* **554**, 548 (2001).
- [38] B. Willems, and V. Kalogera, *ApJ* **603**, L101 (2004).

- [39] L.R. Yungelson, and M. Livio, *ApJ* **528**, 108 (2000).
- [40] A.V. Fedorova, A.V. Tutukov, and L.R. Yungelson, *Astron. Lett.* **30**, 73 (2004).
- [41] R. Napiwotzki, et al., e-lanl: astro-ph/0403595.
- [42] L.R. Yungelson, in *White Dwarfs: Galactic and Cosmological Probes*, Edited by E.M. Sion, H.L. Shipman, and S. Vennes, Kluwer Academic Publishers, in press; e-lanl: astro-ph/0409677.
- [43] G. Nelemans, L.R. Yungelson, and S.F. Portegies Zwart, *A&A* **375**, 890 (2001).
- [44] O. Pols, and M. Marinus, *A&A* **288**, 475 (1994).
- [45] H. Bethe, and G.E. Brown, *ApJ* **506**, 780 (1998).
- [46] S.F. Portegies-Zwart, and L.R. Yungelson, *A&A* **332**, 173 (1998).
- [47] J. Bloom, S. Sigurdsson, and O. Pols, *MNRAS* **305**, 763 (1999).
- [48] K. Belczynski, and T. Bulik, *A&A* **346**, 91 (1999).
- [49] K. Belczynski, T. Bulik, and W. Kluźniak, *ApJ* **567**, L63 (2002).
- [50] A.V. Tutukov, and L.R. Yungelson, *Astron. Rep.* **46**, 667 (2002).
- [51] U. Kolb, *ASP Conference Series* **137**, 190 (1998).
- [52] U. Kolb, A.R. King, and H. Ritter, *MNRAS* **298**, L29 (1998).
- [53] S.B. Howell, L.A. Nelson, and S. Rappaport, *ApJ* **550**, 897 (2001).
- [54] B. Willems, and U. Kolb, *A&A* (2004) in press; e-lanl: astro-ph/0403090.
- [55] U. Kolb, and B. Willems, *Rev. Mex. A&A* **20**, 101 (2004).
- [56] I. Iben, A.V. Tutukov, and L.R. Yungelson, *ApJ* **475**, 291 (1997).
- [57] G. Nelemans, and C.A. Tout, *Rev. Mex. A&A* **20**, 39 (2004).
- [58] M. Politano, *AAS* **199**, 6208 (2001).
- [59] G. Nelemans, L.R. Yungelson, S.F. Portegies Zwart, and F. Verbunt, *A&A* **365**, 491 (2001).
- [60] F. Manucci et al., e-lanl: 0411450.

- [61] P.C. Joss, and S.A. Rappaport, ARA&A **22**, 537 (1984).
- [62] F. Verbunt, ARA&A **31**, 93 (1993).
- [63] Y. Tanaka, and N. Shibazaki, ARA&A **34**, 607 (1996).
- [64] M. van der Kliss, ARA&A **38**, 717 (2000).
- [65] J. Swank, and C. Markwardt, ASP Conf. **251**, 94 (2001).
- [66] H.-J. Grimm, M. Gilfanov, and R. Sunyaev, Memorie Della Società Astronomica Italiana **73**, 1053 (2002).
- [67] W.W. Dalton, and C.L. Sarazin, ApJ **440**, 280 (1995).
- [68] J.L. Terman, R.E. Taam, and C.O. Savage, MNRAS **293**, 113 (1998).
- [69] I. Iben, A.V. Tutukov, and L.R. Yungelson, ApJS **100**, 217 (1995)
I. Iben, A.V. Tutukov, and L.R. Yungelson, ApJS **100**, 233 (1995)
- [70] V.M. Lipunov, L.M. Ozernoy, S.B. Popov, K.A. Postnov, and M.E. Prokhorov, ApJ **466**, 234 (1996).
- [71] W. Kinwah, PASA **18**, 443 (2001).
- [72] Ph. Podsiadlowski, S. Rappaport, and E. Pfahl, in *The influence of binaries on stellar population studies*, Kluwer Academic Publishers, Dordrecht, Astrophysics and space science library, v.264, p.355 (2001).
- [73] E. Pfahl, S. Rappaport, and Ph. Podsiadlowski, ApJ **597**, 1036 (2003).
- [74] A. Possenti, M. Colpi, N. D'Amico, and L. Burderi, ApJ **497**, L97 (1998).
- [75] B. Willems, and U. Kolb, MNRAS **337**, 1004 (2002).
- [76] P.P. Eggleton, and F. Verbunt, MNRAS **220**, P13 (1986).
- [77] R. Mardling, and S. Aarseth, in *The Dynamics of Small Bodies in the Solar System, A Major Key to Solar System Studies*, Edited by B.A. Steves and A.E. Roy, Kluwer Academic Publishers, p.385 (1999).
- [78] A.G. Kuranov, K.A. Postnov, and M.E. Prokhorov, Astron. Rep. **45**, 620 (2001).
- [79] E.J.M. Colbert, and A. Ptak, ApJS **143**, 25 (2002).

- [80] R. Mushotzky, e-lanl: astro-ph/0411040.
- [81] G. Fabbiano, *ARA&A* **27**, 87 (1989).
- [82] E.J.M. Colbert, and R.F. Mushotzky, *ApJ* **519**, 89 (1999).
- [83] M. Coleman Miller, and E.J.M. Colbert, *Int.J.Mod.Phys.* **D13**, 1 (2004).
- [84] A.R. King, et al., *ApJ* **552**, L109 (2001).
- [85] K.A. Postnov, *Astron. Lett.* **29**, 372 (2003).
- [86] R. Soria, M. Cropper, and C. Motch, e-lanl: astro-ph/0409130.
- [87] Ph. Podsiadlowski, S. Rappaport, Z. Han, *MNRAS* **341**, 385 (2003).
- [88] S. Chandrasekhar, *Physical Review Letters* **24**, 611 (1970).
- [89] J.L. Friedman, and B.F. Schutz, *ApJ* **222**, 881 (1978).
- [90] K.D. Kokkotas, and N. Stergioulas, *A&A* **341**, 110 (1999).
- [91] N. Andersson, K.D. Kokkotas, and N. Stergioulas, *ApJ* **516**, 307 (1999).
- [92] V.N. Mironovskij, *Astron. Zhurn.* **42**, 977 (1965).
- [93] V.M. Lipunov, K.A. Postnov, and M.E. Prokhorov, *A&A* **176**, L1 (1987).
- [94] L.P. Grishchuk, V.M. Lipunov, K.A. Postnov, M.E. Prokhorov, and S.B. Sathyaprakash, *Phys.Usp.*, **44**, 1 (2001).
- [95] G. Nelemans, *AIPC* **686**, 263 (2003).
- [96] A. Cooray, *MNRAS* **354**, 25 (2004).
- [97] H. Jorgensen, V.M. Lipunov, I.E. Panchenko, K.A. Postnov, and M.E. Prokhorov, *Ap&SS* **231**, 389 (1995).
- [98] E.S. Phinney, *ApJ* **380**, L17 (1991).
- [99] E.P.J. van den Heuvel, and D.R. Lorimer, *MNRAS* **283**, L37 (1996).
- [100] M. Burgay, N. D’Amico, A. Possenti, et al., *Nature* **426**, 531 (2003).
- [101] V. Kalogera, et al., *ApJ* **601**, L179 (2004).
- [102] V.M. Lipunov, e-lanl: astro-ph/0406502.

- [103] V.M. Lipunov, K.A. Postnov, and M.E. Prokhorov, *A&A* **310**, 489 (1996).
- [104] B. Willems, et al., e-lanl: astro-ph/0411423.
- [105] P. Hut, J.N. Bahcall, *ApJ* **268**, 319 (1983).
- [106] D.C. Heggie, and P. Hut, *ApJS* **85**, 347 (1993).
- [107] F.A. Rasio, S. McMillan, and P. Steve, *ApJ* **438**, L33 (1995).
- [108] J.M. Fregeau, P. Cheung, S.F. Portegies Zwart, and F.A. Rasio, *MNRAS* **352**, 1 (2004).
- [109] S.J. Aarseth, *Cel. Mech. & Dyn. Astron.* **73**, 127 (1999).
- [110] J.R.. Hurley, C.A. Tout, S.J. Aarseth, and O.R. Pols, *MNRAS* **323**, 630 (2001).
- [111] A. Gualandris, S.F. Portegies Zwart, and P.P. Eggleton, *MNRAS* **350**, 615 1 (2004).
- [112] S.B. Popov, and M.E. Prokhorov, *Trudy GAISH (in Russian)* **72**, 1 (2003).
- [113] T. Regimbau, and J.A. de Freitas Pacheco, *A&A* **374**, 181 (2001).
- [114] P.L. Gonthier, R. Van Guilder, and A.K. Harding, *ApJ* **604**, 775 (2004).
- [115] V.S. Beskin, and S.A. Eliseeva, *PAZh* in press (2005).
- [116] O. Blaes, and M. Rajagopal, *ApJ* **381**, 210 (1991).
- [117] R.A. Manning, R.D. Jeffries, and A.P. Willmore, *MNRAS* **278**, 577 (1996).
- [118] A.G. Lyne, and D.R. Lorimer, *Nature* **369**, 127 (1994).
- [119] Z. Arzoumanian, D.F. Chernoff, and J.M. Cordes, *ApJ* **568**, 289 (2002).
- [120] S.B. Popov, M. Colpi, A. Treves, R. Turolla, V.M. Lipunov, and M.E. Prokhorov, *ApJ* **530**, 896 (2000).
- [121] N.R. Ikhsanov, *A&A* **399**, 1147 (2003).
- [122] S.B. Popov, M. Colpi, M.E. Prokhorov, A. Treves, and R. Turolla, *ApJ* **544**, L53 (2000).
- [123] S.B. Popov, M.E. Prokhorov, M. Colpi, A. Treves, and R. Turolla, *A&A*, **406**, 111 (2003).

- [124] S.B. Popov, M. Colpi, M.E. Prokhorov, A. Treves, and R. Turolla, Ap&SS in press, e-lanl: astro-ph/0305599.
- [125] S.B. Popov, H. Grigorian, R. Turolla, and D. Blaschke, e-lanl: astro-ph/0411618.
- [126] S.B. Popov, A. Treves, and R. Turolla, in proceedings of the *4th AGILE workshop*, Edited by M. Tavani, A. Pellizzoni, and S. Vercellone. Aracne Editrice, p. 183, e-lanl: astro-ph/0310416 (2004).
- [127] F. Zhang, Z. Han, L. Li, and J.R. Hurley, MNRAS **350**, 710 (2004).
- [128] C. Leitherer, in *The Evolution of Starbursts*, Edited by S. Huettemeister and E. Manthey, in press, e-lanl: astro-ph/0409407.
- [129] J.M. Mas-Hesse, and M. Cerviño, in *Wolf-Rayet Phenomena in Massive Stars and Starburst Galaxies*, IAU Symp. 193, Edited by K.A. van der Hucht, G. Koenigsberger, and P.R.J. Eenens, p. 550 (1999).
- [130] F. Zhang, Z. Han, L. Li, and J.R. Hurley, A&A **415**, 117 (2004).
- [131] E. De Donder, and D. Vanbeveren New Astron.Rev. **48** 861 (2004).
- [132] C. Leitherer, e-lanl: astro-ph/0411345.
- [133] E.F. Bell, and R.S. de Jong, ApJ **550**, 212 (2001).
- [134] F. Pompilio, F. La Franca, and G. Matt, A&A **353**, 440 (2000).
- [135] P. Madau, G. Ghisellini, and A.C. Fabian, MNRAS **270**, 17 (1994).
- [136] R.R.J. Antonucci, ARA&A **31**, 473 (1993).
- [137] R. Gilli, M. Salvati, and G. Hasinger, A&A **366**, 407 (2001).
- [138] P. Gandhi, A.C. Fabian, MNRAS **339**, 1095 (2003).
- [139] A.V. Halevin, e-lanl: astro-ph/0309176.